

Thank you for requesting Medicis before and after photos of lip augmentation with *Restylane*®.
 The images below represent the photos currently available for your selection.
 Please identify which sets of photos you would like to receive and we will send copies electronically.

Restylane—Lip Before and After Images (Print and Web)



Lips before treatment



After 3.4 mL of *Restylane*®
 (including initial and touch-up treatments)
 8 weeks post-treatment. Individual results may vary.

RES 11-162B 02/28/13 Customer Service_B&A



Lips before treatment



After 2 mL of *Restylane*®
 (including initial and touch-up treatments)
 Immediately following treatment. Individual results may vary.

RES 11-162E 02/28/13 Customer Service_B&A



Lips before treatment



After 2.7 mL of *Restylane*®
 (including initial and touch-up treatments)
 8 weeks post-treatment. Individual results may vary.

RES 11-162C 02/28/13 Customer Service_B&A



Before treatment
 (Lips and oral commissures)



After 1.3 mL of *Restylane*®
 (including initial and touch-up treatments)
 Immediately following treatment. Individual results may vary.

RES 11-162F 02/28/13 Customer Service_B&A



Lips before treatment



After 2.6 mL of *Restylane*®
 (including initial and touch-up treatments)
 8 weeks post-treatment. Individual results may vary.

RES 11-162D 02/28/13 Customer Service_B&A



Before treatment
 (Lips, oral commissures, marionette lines, perioral rhytides)



After 3.5 mL of *Restylane*®
 (including initial and touch-up treatments)
 Immediately following treatment. Individual results may vary.

RES 11-162G 02/28/13 Customer Service_B&A

Each lip before & after set includes Important Safety Considerations for *Restylane*.



See Important Safety Information for *Restylane* on the reverse side.

Restylane is indicated for submucosal implantation for lip augmentation in patients over 21 years and for mid-to-deep dermal implantation for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds.

Important Safety Information for *Restylane*

Restylane contains traces of gram-positive bacterial protein and is contraindicated for patients with allergies to such material or for patients with severe allergies that have required in-hospital treatment. *Restylane* should not be used by patients with bleeding disorders. *Restylane* should not be injected anywhere except the dermis or lip submucosa.

Use of *Restylane* at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. The most commonly observed side effects are swelling, redness, pain, bruising, and tenderness at the injection site. These are typically mild in severity and resolve in less than 7 days in nasolabial folds and less than 14 days in lips. The incidence of swelling may be higher in patients under 36 years, and the incidence of bruising may be higher in patients over 35 years. Serious but rare side effects include delayed onset infections, recurrence of herpetic eruptions, and superficial necrosis and scarring at the injection site. Do not implant into blood vessels. Use with caution in patients recently treated with anticoagulant or platelet inhibitors to avoid bleeding and bruising. Safety has not been established for use during pregnancy, when breastfeeding, or in patients under 18 years for nasolabial folds.

Treatment volume should be limited to 6.0 mL in wrinkles and folds, such as nasolabial folds, and limited to 1.5 mL per lip, as higher volume significantly increases moderate and severe injection site reactions. The safety or effectiveness of treatment in areas other than nasolabial folds and lips has not been established in controlled clinical studies.

Restylane is available only through a licensed practitioner. Complete product and safety information is available at www.RestylaneUSA.com.



Restylane is a registered trademark of HA North American Sales AB.
RES 11-162A 02/28/13

Restylane[®]
INJECTABLE GEL

BY MEDICIS AESTHETICS